

nodular masses is also involved over the greater part of the organ. On the anterior wall, about 4 cm. from the cardia, is a perforation the size of a three-penny bit. For 1 cm. about it the mucosa is ulcerated. No other spot of ulceration exists.

The left supra-renal capsule is enlarged, and partially involved in cancerous disease. About it also are several small firm nodules.

Nothing special in the other organs.

(b.)—*Cancer of Stomach—Extensive Ulceration of Anterior Wall—Left Lobe of Liver projecting into the Organ.*

Sarah W., æt. 43. Ill for six months with all the ordinary symptoms of cancer of stomach. Moderate emaciation. General œdema—slight in trunk.

In *abdomen*, parietal peritoneum adherent to omentum in epigastric region. About two quarts of turbid fluid removed. Omentum retracted and puckered. Left lobe of liver and stomach firmly united together. *Stomach* of average size. Orifices free. On slitting it open along the greater curvature, a large ulcerated surface is seen on the anterior wall and lesser curvature, nearer the pylorus than the cardia, and about the size of the palm of the hand in extent. On pouring water upon it, part of the left lobe of the liver is seen to project through the centre, while the peripheral parts are in a soft, sloughy state. The walls at the margins of the ulcer are infiltrated and thickened, but not to a very great extent.

No secondary masses in other organs.

(c.)—*Cancer of Stomach—Flattened Mass, not Ulcerated—Secondary Masses in Mesenteric Glands, Pancreas and Gall-bladder.*

Mary J., æt. 60. Fairly well-nourished. Admitted into